



Funded by the
European Union



Tensions in Cosmology: Are we approaching New Physics?

Emmanuel N. Saridakis

National Observatory of Athens

Dragovich-80 May 2025



- The **history** of **Astronomy, Cosmology** and **Gravity** is a **history of tensions** between **theoretical predictions** and **observations**
- **Astrophysical cosmology** has become a **precision science** with an incredibly **huge** amount of data
- **New Tensions appear.**
Are we approaching New Physics?

Aristotle - 350 BC

- According to Aristotle heavier bodies fall faster.
- Bodies fall in order to come back to their “initial state”.




Schema huius præmissæ diuisionis Sphærarum.



China, India, Persia, Arabic Observation

- Observations in China, India, Persia, e.g. in Maragha in 11th century, started putting into doubt Earth's non-motion, however not geocentrism.





Dunhuang map from the Tang dynasty 710 AD




Brahe, Kepler- 1600

- Heliocentrism, elliptical Orbits



Galileo - 1600

- Bodies fall with the same speed, **independently** from their **weight**.





Newton - 1700

- Law of Universal Gravitation:

All bodies (either apples or planets) **attract mutually**.


First time that **gravity is related to astronomy**



$$F_1 = F_2 = G \frac{m_1 \times m_2}{r^2}$$


Mercury perihelion - 1859

- The true orbits of planets, even if seen from the SUN are not ellipses. They are rather curves of this type:*




This angle is the perihelion advance, predicted by G.R.

Michelson–Morley experiment - 1887



General Relativity

- Einstein 1915: **General Relativity**



energy-momentum source of spacetime
Curvature

$$S = \frac{1}{16\pi G} \int d^4x \sqrt{-g} [R - 2\Lambda] + \int d^4x L_m(g_{\mu\nu}, \psi)$$

$$\Rightarrow R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + g_{\mu\nu} \Lambda = 8\pi G T_{\mu\nu}$$

$$T^{\mu\nu} \equiv \frac{2}{\sqrt{-g}} \frac{\delta L_m}{\delta g_{\mu\nu}}$$







with

Modified Gravity before General Relativity

- Modifications to **Newton's Law**
- **Inverse Cube Law**.
- **Extended Inverse-Square Law** (Simon Newcomb - 1880's)
- **Lord Kelvin** - theory of everything (end of 19th century)
- **Hendrik Lorentz**: gravity on the basis of his ether theory and Maxwell's equations. (1900)
- **Nordström's theory of gravitation** (1912 and 1913)
- **Einstein's scalar theory of gravity** (1913)

Summary of 20th century Observations

The Universe history:



Standard Model of Cosmology

Λ CDM Paradigm + Inflation

$$H(t)^2 + \frac{k}{a(t)^2} = \frac{8\pi G}{3} [\rho_{dm}(t) + \rho_b(t) + \rho_r(t)] + \frac{\Lambda}{3}$$

$$w_{\Lambda} \equiv \frac{p_{\Lambda}}{\rho_{\Lambda}} = -1$$

$$\dot{H}(t) - \frac{k}{a(t)^2} = -4\pi G [\rho_{dm}(t) + p_{dm}(t) + \rho_b(t) + p_b(t) + \rho_r(t) + p_r(t)]$$

Λ CDM concordance model is almost perfect!

- Describes the thermal history of the Universe at the background level
- Epochs of inflation, radiation, matter, late-time acceleration

Cosmology-background

- Homogeneity and isotro $ds^2 = -dt^2 + a^2(t) \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right)$
- Background evolution (Friedmann equations) in flat space

$$H^2 = \frac{8\pi G}{3} (\rho_m + \rho_{DE})$$

$$\dot{H} = -4\pi G (\rho_m + p_m + \rho_{DE} + p_{DE}),$$

(the effective DE sector can be either Λ or any possible modification)

- One must obtain a $H(z)$ and $\Omega_m(z)$ and $w_{DE}(z)$ in agreement with observations (SNIa, BAO, CMB shift parameter, $H(z)$ etc)

Cosmology-perturbations

- **Perturbation evolution** $\ddot{\delta} + 2H\dot{\delta} - 4\pi G_{\text{eff}} \rho \delta \approx 0$ where $\delta \equiv \delta\rho/\rho$
where $G_{\text{eff}}(z, k)$ is the **effective Newton's constant**, given by

$$\nabla^2 \phi \approx 4\pi G_{\text{eff}} \rho \delta,$$

under the scalar **metric perturbation** $ds^2 = -(1 + 2\phi)dt^2 + a^2(1 - 2\psi)d\vec{x}^2$


- Hence: $\delta'' + \left(\frac{(H^2)'}{2H^2} - \frac{1}{1+z} \right) \delta' \approx \frac{3}{2}(1+z) \frac{H_0^2}{H^2} \frac{G_{\text{eff}}(z, k)}{G_N} \Omega_{0m} \delta$

with $f(a) = \frac{d \ln \delta}{d \ln a}$ the **growth rate**, with $f(a) = \Omega_m(a)^{\gamma(a)}$ and $\Omega_m(a) \equiv \frac{\Omega_{0m} a^{-3}}{H(a)^2/H_0^2}$


- One can define the **observable**: $f\sigma_8(a) \equiv f(a) \cdot \sigma(a) = \frac{\sigma_8}{\delta(1)} a \delta'(a)$

with $\sigma(a) = \sigma_8 \frac{\delta(a)}{\delta(1)}$ the z-dependent rms fluctuations of the linear density field within spheres of radius $R = 8h^{-1}\text{Mpc}$, and σ_8 its value today.





Matter Density Fluctuation Power Spectrum



A different convention:
plot $P(k)k^3$



Cosmology in the 21st century



Issues of Λ CDM Paradigm

- 1) General Relativity is non-renormalizable. It cannot get quantized.
- 2) The cosmological-constant problem.
- 3) How to describe primordial universe (inflation)
- 4) Physics of Dark Matter
- 5) A huge amount of accumulating data suggest possible tensions:


H_0 , $f\sigma_8$

Challenges for Λ CDM Beyond H_0 and S_8

- A. The A_{lens} Anomaly in the CMB Angular Power Spectrum
- B. Hints for a Closed Universe from *Planck* Data
- C. Large-Angular-Scale Anomalies in the CMB Temperature and Polarization
 - 1. The Lack of Large-Angle CMB Temperature Correlations
 - 2. Hemispherical Power Asymmetry
 - 3. Quadrupole and Octopole Anomalies
 - 4. Point-Parity Anomaly
 - 5. Variation in Cosmological Parameters Over the Sky
 - 6. The Cold Spot
 - 7. Explaining the Large-Angle Anomalies
 - 8. Predictions and Future Testability
 - 9. Summary
- D. Abnormal Oscillations of Best Fit Parameter Values
- E. Anomalously Strong ISW Effect
- F. Cosmic Dipoles
 - 1. The α Dipole
 - 2. Galaxy Cluster Anisotropies and Anomalous Bulk Flows
 - 3. Radio Galaxy Cosmic Dipole
 - 4. QSO Cosmic Dipole and Polarisation Alignments
 - 5. Dipole in SNIa
 - 6. Emergent Dipole in H_0
 - 7. CMB Dipole: Intrinsic Versus Kinematic?
- G. The Ly- α Forest BAO and CMB Anomalies
 - 1. The Ly- α Forest BAO Anomaly
 - 2. Ly- α -*Planck* 2018 Tension in n_s - Ω_m
- H. Parity Violating Rotation of CMB Linear Polarization
 - 1. The Lithium Problem
- J. Quasars Hubble Diagram Tension with Planck- Λ CDM
- K. Oscillating Force Signals in Short Range Gravity Experiments
- L. Λ CDM and the Dark Matter Phenomenon at Galactic Scales

[L. Perivolaropoulos , F. Scara, New Astron.Rev (2022), 2105.05208 [astro-ph.CO]]

Can General Relativity be quantized?



COSMOLOGICAL CONSTANT PROBLEM

$$E_n \sim (n + 1/2)h\omega(k)$$

$$\rho_\Lambda(th) \sim M_p^4$$

$$\rho_\Lambda^0 \sim 10^{-120} \rho_\Lambda^{th}$$

H0 tension

- **Tension (5σ !)** between the **data** (direct measurements) and **Planck/ Λ CDM** (indirect measurements). The data indicate **a lack of “gravitational power”**.






Figure 26. The CMB power spectrum as a function of cosmological parameters


[Riess et al, Astrophys.] 826]

Current status

H0 measured / inferred using many techniques




“Cosmology Intertwined” (2022)



[Abdalla et al, JHEAp (2022)] 21

H0 tension

- **Tension** between the **data** (direct measurements) and **Planck/ Λ CDM** (indirect measurements). This tension could be due to **systematics**.
- If not systematics then we may need **changes in Λ CDM** in **early** or **late** time behavior. **5σ** seems to be very serious!




- Change early or late Universe physics. **Higher number** of effective **relativistic species**, **dynamical dark energy**, **non-zero curvature**, etc.
- The data indicate **a lack of “gravitational power”**. **Modified Gravity**.

Restoring cosmological concordance

Is LCDM Wrong?

$$\theta_s = \frac{r_s}{D_A}$$

0.04% precision



$$r_s \propto \int_0^{t_{\text{recom}}} dt \frac{c_s(t)}{\rho(t)}$$

$$D_A \propto \frac{1}{H_0} \int_{t_{\text{recom}}}^{t_{\text{today}}} dt \frac{1}{\rho(t)}$$

How do we increase H_0 ?

Decrease sound horizon (r_s)


Increase integral in angular diameter distance (D_A)

“Early time solutions”

“Late time solutions”

S8 Tension

- Tension** between **direct data** and **Planck/ Λ CDM estimation**. The data indicate **less matter clustering** in structures at intermediate-small cosmological scales.



S8 Tension

TABLE II: A compilation of RSD data that we found published from 2006 since 2018

Index	Dataset	z	$f\sigma_8(z)$	Refs.	Year	Fiducial Cosmology
1	SDSS-LRG	0.35	0.440 ± 0.050	[75]	30 October 2006	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.25, 0, 0.756)[76]$
2	VVDS	0.77	0.490 ± 0.18	[75]	6 October 2009	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.25, 0, 0.78)$
3	2dFGRS	0.17	0.510 ± 0.060	[75]	6 October 2009	$(\Omega_{0m}, \Omega_K) = (0.3, 0, 0.9)$
4	2MRS	0.02	0.314 ± 0.048	[77], [78]	13 November 2010	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.266, 0, 0.65)$
5	SnIa+IRAS	0.02	0.398 ± 0.065	[79], [78]	20 October 2011	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.3, 0, 0.814)$
6	SDSS-LRG-200	0.25	0.3512 ± 0.0583	[80]	9 December 2011	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.276, 0, 0.8)$
7	SDSS-LRG-200	0.37	0.4602 ± 0.0378	[80]	9 December 2011	
8	SDSS-LRG-60	0.25	0.3665 ± 0.0601	[80]	9 December 2011	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.276, 0, 0.8)$
9	SDSS-LRG-60	0.37	0.4031 ± 0.0586	[80]	9 December 2011	
10	WiggleZ	0.44	0.413 ± 0.080	[46]	12 June 2012	$(\Omega_{0m}, h, \sigma_8) = (0.27, 0.71, 0.8)$
11	WiggleZ	0.60	0.390 ± 0.063	[46]	12 June 2012	$C_{ij} = Eq.(3.3)$
12	WiggleZ	0.73	0.437 ± 0.072	[46]	12 June 2012	
13	6dFGS	0.067	0.423 ± 0.055	[81]	4 July 2012	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.27, 0, 0.76)$
14	SDSS-BOSS	0.30	0.407 ± 0.055	[82]	11 August 2012	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.25, 0, 0.804)$
15	SDSS-BOSS	0.40	0.419 ± 0.041	[82]	11 August 2012	
16	SDSS-BOSS	0.50	0.427 ± 0.043	[82]	11 August 2012	
17	SDSS-BOSS	0.60	0.433 ± 0.067	[82]	11 August 2012	
18	Vipers	0.80	0.470 ± 0.080	[83]	9 July 2013	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.25, 0, 0.82)$
19	SDSS-DR7-LRG	0.35	0.429 ± 0.089	[84]	8 August 2013	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.25, 0, 0.809)[85]$
20	GAMA	0.18	0.360 ± 0.090	[86]	22 September 2013	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.27, 0, 0.8)$
21	GAMA	0.38	0.440 ± 0.060	[86]	22 September 2013	
22	BOSS-LOWZ	0.32	0.384 ± 0.095	[87]	17 December 2013	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.274, 0, 0.8)$
23	SDSS DR10 and DR11	0.32	0.48 ± 0.10	[87]	17 December 2013	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.274, 0, 0.8)[88]$
24	SDSS DR10 and DR11	0.57	0.417 ± 0.045	[87]	17 December 2013	
25	SDSS-MGS	0.15	0.490 ± 0.145	[89]	30 January 2015	$(\Omega_{0m}, h, \sigma_8) = (0.31, 0.67, 0.83)$
26	SDSS-veloc	0.10	0.370 ± 0.130	[90]	16 June 2015	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.3, 0, 0.89)[91]$
27	FastSound	1.40	0.482 ± 0.116	[92]	25 November 2015	$(\Omega_{0m}, h, \sigma_8) = (0.27, 0, 0.82)[93]$
28	SDSS-CMASS	0.59	0.488 ± 0.060	[94]	8 July 2016	$(\Omega_{0m}, h, \sigma_8) = (0.307115, 0.6777, 0.8288)$
29	BOSS DR12	0.38	0.497 ± 0.045	[2]	11 July 2016	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.31, 0, 0.8)$
30	BOSS DR12	0.51	0.458 ± 0.038	[2]	11 July 2016	
31	BOSS DR12	0.61	0.436 ± 0.034	[2]	11 July 2016	
32	BOSS DR12	0.38	0.477 ± 0.051	[95]	11 July 2016	$(\Omega_{0m}, h, \sigma_8) = (0.31, 0.676, 0.8)$
33	BOSS DR12	0.51	0.453 ± 0.050	[95]	11 July 2016	
34	BOSS DR12	0.61	0.410 ± 0.044	[95]	11 July 2016	
35	Vipers v7	0.76	0.440 ± 0.040	[55]	26 October 2016	$(\Omega_{0m}, \sigma_8) = (0.308, 0.8149)$
36	Vipers v7	1.05	0.280 ± 0.080	[55]	26 October 2016	
37	BOSS LOWZ	0.32	0.427 ± 0.056	[96]	26 October 2016	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.31, 0, 0.8475)$
38	BOSS CMASS	0.57	0.426 ± 0.029	[96]	26 October 2016	
39	Vipers	0.727	0.296 ± 0.0765	[97]	21 November 2016	$(\Omega_{0m}, \Omega_K, \sigma_8) = (0.31, 0, 0.7)$
40	6dFGS+SnIa	0.02	0.428 ± 0.0465	[98]	29 November 2016	$(\Omega_{0m}, h, \sigma_8) = (0.3, 0.683, 0.8)$
41	Vipers	0.6	0.48 ± 0.12	[99]	16 December 2016	$(\Omega_{0m}, \Omega_b, n_s, \sigma_8) = (0.3, 0.045, 0.96, 0.831)[12]$
42	Vipers	0.86	0.48 ± 0.10	[99]	16 December 2016	
43	Vipers PDR-2	0.60	0.550 ± 0.120	[100]	16 December 2016	$(\Omega_{0m}, \Omega_b, \sigma_8) = (0.3, 0.045, 0.823)$
44	Vipers PDR-2	0.86	0.400 ± 0.110	[100]	16 December 2016	
45	SDSS DR13	0.1	0.48 ± 0.16	[101]	22 December 2016	$(\Omega_{0m}, \sigma_8) = (0.25, 0.89)[91]$
46	2MTF	0.001	0.505 ± 0.085	[102]	16 June 2017	$(\Omega_{0m}, \sigma_8) = (0.3121, 0.815)$
47	Vipers PDR-2	0.85	0.45 ± 0.11	[103]	31 July 2017	$(\Omega_b, \Omega_{0m}, h) = (0.045, 0.30, 0.8)$
48	BOSS DR12	0.31	0.469 ± 0.098	[49]	15 September 2017	$(\Omega_{0m}, h, \sigma_8) = (0.307, 0.6777, 0.8288)$
49	BOSS DR12	0.36	0.474 ± 0.097	[49]	15 September 2017	
50	BOSS DR12	0.40	0.473 ± 0.086	[49]	15 September 2017	
51	BOSS DR12	0.44	0.481 ± 0.076	[49]	15 September 2017	
52	BOSS DR12	0.48	0.482 ± 0.067	[49]	15 September 2017	
53	BOSS DR12	0.52	0.488 ± 0.065	[49]	15 September 2017	
54	BOSS DR12	0.56	0.482 ± 0.067	[49]	15 September 2017	
55	BOSS DR12	0.59	0.481 ± 0.066	[49]	15 September 2017	
56	BOSS DR12	0.64	0.486 ± 0.070	[49]	15 September 2017	
57	SDSS DR7	0.1	0.376 ± 0.038	[104]	12 December 2017	$(\Omega_{0m}, \Omega_b, \sigma_8) = (0.282, 0.046, 0.817)$
58	SDSS-IV	1.52	0.420 ± 0.076	[105]	8 January 2018	$(\Omega_{0m}, \Omega_b h^2, \sigma_8) = (0.26479, 0.02258, 0.8)$
59	SDSS-IV	1.52	0.396 ± 0.079	[106]	8 January 2018	$(\Omega_{0m}, \Omega_b h^2, \sigma_8) = (0.31, 0.022, 0.8225)$
60	SDSS-IV	0.978	0.379 ± 0.176	[107]	9 January 2018	$(\Omega_{0m}, \sigma_8) = (0.31, 0.8)$
61	SDSS-IV	1.23	0.385 ± 0.099	[107]	9 January 2018	
62	SDSS-IV	1.526	0.342 ± 0.070	[107]	9 January 2018	
63	SDSS-IV	1.944	0.364 ± 0.106	[107]	9 January 2018	

- **Model Dependence:** Distance to galaxies is not measured directly, so a cosmological model is assumed in order to infer distances (Λ CDM with different parameters).
- **Double counting:** Some data points correspond to the same sample of galaxies analyzed by different groups/methods etc.

[Kazantzidis, Perivolaropoulos, PRD97]

Tension2 – $f\sigma_8$

- **Tension** between the **data** and **Planck/ Λ CDM**.
- This tension could be due to **systematics**.
- **If not systematics**, the data **less matter clustering** in structures at intermediate-small cosmological scales (expressed as smaller Ω_m at $z < 0.6$, or smaller σ_8 , or $w_{DE} < -1$).
- It could be reconciled by a **mechanism that reduces the rate of clustering** between recombination and today: **Hot Dark Matter**, **Dark Matter that clusters differently** at small scales, or **Modified Gravity**.

Possible Solutions of H0 and S8 tensions

tension $\leq 1\sigma$ “Excellent models”	tension $\leq 2\sigma$ “Good models”	tension $\leq 3\sigma$ “Promising models”
Dark energy in extended parameter spaces [289] Dynamical Dark Energy [309] Metastable Dark Energy [314] PEDE [392, 394] Elaborated Vacuum Metamorphosis [400–402] IDE [314, 636, 637, 639, 652, 657, 661–663] Self-interacting sterile neutrinos [711] Generalized Chaplygin gas model [744] Galileon gravity [876, 882] Power Law Inflation [966] $f(\mathcal{T})$ [818]	Early Dark Energy [235] Phantom Dark Energy [11] Dynamical Dark Energy [11, 281, 309] GEDE [397] Vacuum Metamorphosis [402] IDE [314, 653, 656, 661, 663, 670] Critically Emergent Dark Energy [997] $f(\mathcal{T})$ gravity [814] Über-gravity [59] Reconstructed PPS [978]	Early Dark Energy [229] Decaying Warm DM [474] Neutrino-DM Interaction [506] Interacting dark radiation [517] Self-Interacting Neutrinos [700, 701] IDE [656] Unified Cosmologies [747] Scalar-tensor gravity [856] Modified recombination [986] Super Λ CDM [1007] Coupled Dark Energy [650]
Early Dark Energy [228, 235, 240, 250] Exponential Acoustic Dark Energy [259] Phantom Crossing [315] Late Dark Energy Transition [317] Metastable Dark Energy [314] PEDE [394] Vacuum Metamorphosis [402] Elaborated Vacuum Metamorphosis [401, 402] Sterile Neutrinos [433] Decaying Dark Matter [481] Neutrino-Majoron Interactions [509] IDE [637, 639, 657, 661] DM - Photon Coupling [685] $f(\mathcal{T})$ gravity theory [812] BD- Λ CDM [851] Über-Gravity [59] Galileon Gravity [875] Unimodular Gravity [890] Time Varying Electron Mass [990] Λ CDM [995] Ginzburg-Landau theory [996] Lorentzian Quintessential Inflation [979] Holographic Dark Energy [351]	Early Dark Energy [212, 229, 236, 263] Rock ‘n’ Roll [242] New Early Dark Energy [247] Acoustic Dark Energy [257] Dynamical Dark Energy [309] Running vacuum model [332] Bulk viscous models [340, 341] Holographic Dark Energy [350] Phantom Braneworld DE [378] PEDE [391, 392] Elaborated Vacuum Metamorphosis [401] IDE [659, 670] Interacting Dark Radiation [517] Decaying Dark Matter [471, 474] DM - Photon Coupling [686] Self-interacting sterile neutrinos [711] $f(\mathcal{T})$ gravity theory [817] Über-Gravity [871] VCDM [893] Primordial magnetic fields [992] Early modified gravity [859] Bianchi type I spacetime [999] $f(\mathcal{T})$ [818]	DE in extended parameter spaces [289] Dynamical Dark Energy [281, 309] Holographic Dark Energy [350] Swampland Conjectures [370] MEDE [399] Coupled DM - Dark radiation [534] Decaying Ultralight Scalar [538] BD- Λ CDM [852] Metastable Dark Energy [314] Self-Interacting Neutrinos [700] Dark Neutrino Interactions [716] IDE [634–636, 653, 656, 663, 669] Scalar-tensor gravity [855, 856] Galileon gravity [877, 881] Nonlocal gravity [886] Modified recombination [986] Effective Electron Rest Mass [989] Super Λ CDM [1007] Axi-Higgs [991] Self-Interacting Dark Matter [479] Primordial Black Holes [545]

Possible Solutions of H0 and S8 tensions

Early-Time Alternative Proposed Models

1. Axion Monodromy
2. Early Dark Energy
3. Extra Relativistic Degrees of Freedom
4. Modified Recombination History
5. New Early Dark Energy

Late-Time Alternative Proposed Models

1. Bulk Viscous Models
2. Chameleon Dark Energy
3. Clustering Dark Energy
4. Diffusion Models
5. Dynamical Dark Energy
6. Emergent Dark Energy
7. Graduated Dark Energy - AdS to dS Transition in the Late Universe
8. Holographic Dark Energy
9. Interacting Dark Energy
10. Quintessence Models and their Various Extensions
11. Running Vacuum Models
12. Time-Varying Gravitational Constant
13. Vacuum Metamorphosis

Modified Gravity Models

1. Effective Field Theory Approach to Dark Energy and Modified Gravity
2. $f(T)$ Gravity
3. Horndeski Theory
4. Quantum Conformal Anomaly Effective Theory and Dynamical Vacuum Energy
5. Ultra-Late Time Gravitational Transitions

Beyond the FLRW Framework

1. Cosmological Fitting and Averaging Problems
2. Data Analysis in an Universe with Structure: Accounting for Regional Inhomogeneity and Anisotropy
3. Local Void Scenario

Specific Solutions Assuming FLRW

1. Active and Sterile Neutrinos
2. Cannibal Dark Matter
3. Decaying Dark Matter
4. Dynamical Dark Matter
5. Extended Parameter Spaces Involving A_{lens}
6. Cosmological Scenario with Features in the Primordial Power Spectrum
7. Interacting Dark Matter
8. Quantum Landscape Multiverse
9. Quantum Fisher Cosmology
10. Quartessence
11. Scaling Symmetry and a Mirror Sector
12. Self-Interacting Neutrinos
13. Self-Interacting Sterile Neutrinos
14. Soft Cosmology
15. Two-Body Decaying Cold Dark Matter into Dark Radiation and Warm Dark Matter

**Cosmology Intertwined:
A Review of the Particle Physics, Astrophysics, and Cosmology
Associated with the Cosmological Tensions and Anomalies**

Elcio Abdalla,¹ Guillermo Franco Abellán,² Amin Aboubrahim,³ Adriano Agnello,⁴ Özgür Akarsu,⁵ Yashar Akrami,^{6,7,8,9} George Alestas,¹⁰ Daniel Aloni,¹¹ Luca Amendola,¹² Luis A. Anchordoqui,^{13,14,15} Richard I. Anderson,¹⁶ Nikki Arendse,¹⁷ Marika Asgari,^{18,19} Mario Ballardini,^{20,21,22,23} Vernon Barger,²⁴ Spyros Basilakos,^{25,26} Ronaldo C. Batista,²⁷ Elia S. Battistelli,^{28,29} Richard Battye,³⁰ Micol Benetti,^{31,32} David Benisty,^{33,34,35} Asher Berlin,³⁶ Paolo de Bernardis,^{28,29} Emanuele Berti,³⁷ Bohdan Bilenko,^{38,39} Simon Birrer,⁴⁰ John P. Blakeslee,⁴¹ Kimberly K. Boddy,⁴² Clecio R. Bom,^{43,44} Alexander Bonilla,⁴⁵ Nicola Borghi,^{46,47} François R. Bouchet,⁴⁸ Matteo Braglia,^{49,21} Thomas Buchert,⁵⁰ Elizabeth Buckley-Geer,^{51,52} Erminia Calabrese,⁵³ Robert R. Caldwell,⁵⁴ David Camarena,⁵⁵ Salvatore Capozziello,^{56,31} Stefano Casertano,⁵⁷ Angela Chen,^{58,59} Geoff C.-F. Chen,⁶⁰ Hsin-Yu Chen,⁶¹ Jens Chluba,³⁰ Anton Chudaykin,⁶² Michele Cicoli,^{20,22} Craig J. Copi,⁶ Fred Courbin,¹⁶ Francis-Yan Cyr-Racine,⁶³ Božena Czerny,⁶⁴ Maria Dainotti,^{65,66,67} Guido D'Amico,^{68,69} Anne-Christine Davis,^{33,34} Javier de Cruz Pérez,⁷⁰ Jaume de Haro,⁷¹ Jacques Delabrouille,^{72,73,74,75} Peter B. Denton,⁷⁶ Suhail Dhawan,⁷⁷ Keith R. Dienes,^{78,79} Eleonora Di Valentino,^{80,*} Pu Du,⁸¹ Dominique Eckert,⁸² Celia Escamilla-Rivera,⁸³ Agnès Ferté,⁸⁴ Fabio Finelli,^{85,22} Pablo Fosalba,^{86,87} Wendy L. Freedman,⁵² Noemi Frusciante,⁸⁸ Enrique Gaztañaga,^{86,87} William Giarè,^{89,29} Elena Giusarma,⁹⁰ Adrià Gómez-Valent,⁹¹ Will Handley,^{92,93} Ian Harrison,⁹⁴ Luke Hart,³⁰ Dhiraj Kumar Hazra,⁹⁵ Alan Heavens,⁷ Asta Heinesen,⁵⁰ Hendrik Hildebrandt,⁹⁶ J. Colin Hill,^{97,98} Natalie B. Hogg,⁹⁹ Daniel E. Holz,^{52,100,101} Deanna C. Hooper,¹⁰² Nikoo Hosseini-njad,¹⁰³ Dragan Huterer,^{104,105} Mustapha Ishak,¹⁰⁶ Mikhail M. Ivanov,¹⁰⁷ Andrew H. Jaffe,⁷ In Sung Jang,⁵² Karsten Jedamzik,¹⁰⁸ Raul Jimenez,^{109,110} Melissa Joseph,¹¹ Shahab Joudaki,^{111,112} Marc Kamionkowski,³⁷ Tanvi Karwal,¹¹³ Lavrentios Kazantzidis,¹⁰ Ryan E. Keeley,¹¹⁴ Michael Klaser,³ Eiichi Komatsu,^{115,116} Léon V.E. Koopmans,¹¹⁷ Suresh Kumar,¹¹⁸ Luca Lamagna,^{28,29} Ruth Lazkoz,¹¹⁹ Chung-Chi Lee,¹²⁰ Julien Lesgourgues,¹²¹ Jackson Levi Said,^{122,123} Tiffany R. Lewis,¹²⁴ Benjamin L'Huilier,¹²⁵ Matteo Lucca,¹²⁶ Roy Maartens,^{23,127,128} Lucas M. Macri,¹²⁹ Danny Marfatia,¹³⁰ Valerio Marra,^{131,132,133} Carlos J. A. P. Martins,^{134,135} Silvia Masi,^{28,29} Sabino Matarrese,^{136,137,138,139} Arindam Mazumdar,¹⁴⁰ Alessandro Melchiorri,^{28,29} Olga Mena,¹⁴¹ Laura Mersini-Houghton,¹⁴² James Mertens,¹⁴³ Dinko Milaković,^{133,132,144} Yuto Minami,¹⁴⁵ Vivian Miranda,¹⁴⁶ Cristian Moreno-Pulido,¹⁴⁷ Michele Moresco,^{46,47} David F. Mota,¹⁴⁸ Emil Mottola,⁶³ Simone Mozzon,¹⁴⁹ Jessica Muir,¹⁵⁰ Ankan Mukherjee,¹⁵¹ Suvodip Mukherjee,¹⁵⁰ Pavel Naselsky,¹⁵² Pran Nath,¹⁵³ Savvas Nesseris,⁹⁹ Florian Niedermann,¹⁵⁴ Alessio Notari,¹⁵⁵ Rafael C. Nunes,¹⁵⁶ Eoin Ó Colgáin,^{157,158} Kayla A. Owens,⁵² Emre Özülker,⁵ Francesco Pace,^{159,160} Andronikos Paliathanasis,^{161,162} Antonella Palmese,¹⁶³ Supriya Pan,¹⁶⁴ Daniela Paoletti,^{85,22} Santiago E. Perez Bergliaffa,¹⁶⁵ Leandros Perivolaropoulos,¹⁰ Dominic W. Pesce,^{166,167} Valeria Pettorino,¹⁶⁸ Oliver H. E. Philcox,^{169,107} Levon Pogossian,¹⁷⁰ Vivian Poulin,² Gaspard Poulot,⁸⁰ Marco Raveri,¹⁷¹ Mark J. Reid,¹⁷² Fabrizio Renzi,¹⁷³ Adam G. Riess,³⁷ Vivian I. Sabla,⁵⁴ Paolo Salucci,^{174,175} Vincenzo Salzano,¹⁷⁶ Emmanuel N. Saridakis,^{26,75,177} Bangalore S. Sathyaprakash,^{178,179,94} Martin Schmaltz,¹¹ Nils Schöneberg,¹⁸⁰ Dan Scolnic,¹⁸¹ Anjan A. Sen,^{182,183} Neelima Sehgal,¹⁸⁴ Arman Shafieloo,¹⁸⁵ M.M. Sheikh-Jabbari,¹⁸⁶ Joseph Silk,¹⁸⁷ Alessandra Silvestri,¹⁷³ Foteini Slara,¹⁰ Martin S. Sloth,¹⁸⁸ Marcelle Soares-Santos,⁵⁸ Joan Solà Peracaula,¹⁴⁷ Yu-Yang Songsheng,⁸¹ Jorge F. Soriano,^{13,14} Denitsa Staicova,¹⁸⁹ Glenn D. Starkman,^{6,7} István Szapudi,¹⁹⁰ Elsa M. Teixeira,⁸⁰ Brooks Thomas,¹⁹¹ Tommaso Treu,⁶⁰ Emery Trott,⁵⁸ Carsten van de Bruck,⁸⁰ J. Alberto Vazquez,¹⁹² Licia Verde,^{193,194} Luca Visinelli,¹⁹⁵ Deng Wang,¹⁹⁶ Jian-Min Wang,⁸¹ Shao-Jiang Wang,¹⁹⁷ Richard Watkins,¹⁹⁸ Scott Watson,¹⁹⁹ John K. Webb,¹²⁰ Neal Weiner,²⁰⁰ Amanda Weltman,²⁰¹ Samuel J. Witte,²⁰² Radosław Woźtak,⁴ Anil Kumar Yadav,²⁰³ Weiqiang Yang,²⁰⁴ Gong-Bo Zhao,^{205,206} and Miguel Zumalacárregui²⁰⁷

¹Instituto de Física, Universidade de São Paulo - C.P. 66318, CEP: 05315-970, São Paulo, Brazil

²Laboratoire Univers Et Particules de Montpellier (LUPM), Université de Montpellier (UMR-5299)

Efficient model independent requirements to solve the tensions

- In general, to avoid the H_0 tension one needs a positive correction to the first Friedmann equation at late times that could yield an increase in H_0 compared to the Λ CDM scenario.

Efficient model independent requirements to solve the tensions

- For the σ_8 tension, we recall that in any cosmological model, at sub-Hubble scales and through matter epoch, the equation that governs the evolution of matter perturbations in the linear regime is

$$\ddot{\delta} + 2H\dot{\delta} = 4\pi G_{\text{eff}}\rho_m\delta, \quad (1)$$

where G_{eff} is the effective gravitational coupling given by a generalized Poisson equation.

- Solving for $\delta(a)$ provides the observable quantity $f\sigma_8(a)$, following the definitions $f(a) \equiv d \ln \delta(a)/d \ln a$ and $\sigma(a) = \sigma_8 \delta(1)/\delta(a=1)$. Hence, alleviation of the σ_8 tension may be obtained if G_{eff} becomes smaller than G_N during the growth of matter perturbations and/or if the “friction” term in (1) increases.

General Relativity

Assumptions and Considerations


$$S = \frac{1}{16\pi G} \int d^4x \sqrt{-g} [R - 2\Lambda] + \int d^4x L_m(g_{\mu\nu}, \psi)$$

- Diffeomorphism invariance
- Spacetime dimensionality=4
- **Geometry=Curvature** (connection=Levi Civita)
- Linear in Ricci scalar
- **Metric compatibility** (zero non-metricity)
- Minimal matter coupling
- Equivalence principle
- Lorentz invariance
- Locality


Standard Model vs General Relativity Lagrangians

$$\begin{aligned}
 & -\frac{1}{2}\partial_\nu g_\mu^a \partial_\nu g_\mu^a - g_s f^{abc} \partial_\mu g_\nu^a g_\mu^b g_\nu^c - \frac{1}{4}g_s^2 f^{abc} f^{ade} g_\mu^b g_\nu^c g_\mu^d g_\nu^e + \\
 & \frac{1}{2}ig_s^2(\bar{q}_i^\sigma \gamma^\mu q_j^\sigma)g_\mu^a + \bar{G}^a \partial^2 G^a + g_s f^{abc} \partial_\mu \bar{G}^a G^b g_\mu^c - \partial_\nu W_\mu^+ \partial_\nu W_\mu^- - \\
 & M^2 W_\mu^+ W_\mu^- - \frac{1}{2}\partial_\nu Z_\mu^0 \partial_\nu Z_\mu^0 - \frac{1}{2c_w^2}M^2 Z_\mu^0 Z_\mu^0 - \frac{1}{2}\partial_\mu A_\nu \partial_\mu A_\nu - \frac{1}{2}\partial_\mu H \partial_\mu H - \\
 & \frac{1}{2}m_h^2 H^2 - \partial_\mu \phi^+ \partial_\mu \phi^- - M^2 \phi^+ \phi^- - \frac{1}{2}\partial_\mu \phi^0 \partial_\mu \phi^0 - \frac{1}{2c_w^2}M\phi^0 \phi^0 - \beta_h \left[\frac{2M^2}{g^2} + \right. \\
 & \left. \frac{2M}{g}H + \frac{1}{2}(H^2 + \phi^0 \phi^0 + 2\phi^+ \phi^-) \right] + \frac{2M^4}{g^2}\alpha_h - ig_{cw}[\partial_\nu Z_\mu^0(W_\mu^+ W_\nu^- - \\
 & W_\nu^+ W_\mu^-) - Z_\nu^0(W_\mu^+ \partial_\nu W_\mu^- - W_\mu^+ \partial_\nu W_\mu^-) + Z_\mu^0(W_\nu^+ \partial_\nu W_\mu^- - \\
 & W_\nu^+ \partial_\nu W_\mu^-)] - ig_{sw}[\partial_\nu A_\mu(W_\mu^+ W_\nu^- - W_\nu^+ W_\mu^-) - A_\nu(W_\mu^+ \partial_\nu W_\mu^- - \\
 & W_\mu^+ \partial_\nu W_\mu^-) + A_\mu(W_\nu^+ \partial_\nu W_\mu^- - W_\nu^+ \partial_\nu W_\mu^-)] - \frac{1}{2}g^2 W_\mu^+ W_\mu^- W_\nu^+ W_\nu^- + \\
 & \frac{1}{2}g^2 W_\mu^+ W_\nu^- W_\mu^- W_\nu^- + g^2 c_w^2(Z_\mu^0 W_\mu^+ Z_\nu^0 W_\nu^- - Z_\mu^0 W_\nu^+ Z_\nu^0 W_\mu^-) + \\
 & g^2 s_w^2(A_\mu W_\mu^+ A_\nu W_\nu^- - A_\mu A_\nu W_\mu^+ W_\nu^-) + g^2 s_w c_w[A_\mu Z_\nu^0(W_\mu^+ W_\nu^- - \\
 & W_\nu^+ W_\mu^-) - 2A_\mu Z_\mu^0 W_\nu^+ W_\nu^-] - g\alpha[H^3 + H\phi^0 \phi^0 + 2H\phi^+ \phi^-] - \\
 & \frac{1}{8}g^2 \alpha_h[H^4 + (\phi^0)^4 + 4(\phi^+ \phi^-)^2 + 4(\phi^0)^2 \phi^+ \phi^- + 4H^2 \phi^+ \phi^- + 2(\phi^0)^2 H^2] - \\
 & gMW_\mu^+ W_\mu^- H - \frac{1}{2}g\frac{M}{c_w^2}Z_\mu^0 Z_\mu^0 H - \frac{1}{2}ig[W_\mu^+(\phi^0 \partial_\mu \phi^- - \phi^- \partial_\mu \phi^0) - \\
 & W_\mu^-(\phi^0 \partial_\mu \phi^+ - \phi^+ \partial_\mu \phi^0)] + \frac{1}{2}g[W_\mu^+(H\partial_\mu \phi^- - \phi^- \partial_\mu H) - W_\mu^-(H\partial_\mu \phi^+ - \\
 & \phi^+ \partial_\mu H)] + \frac{1}{2}g\frac{1}{c_w}(Z_\mu^0(H\partial_\mu \phi^0 - \phi^0 \partial_\mu H) - ig\frac{s_w}{c_w}MZ_\mu^0(W_\mu^+ \phi^- - W_\mu^- \phi^+) + \\
 & ig_{sw}MA_\mu(W_\mu^+ \phi^- - W_\mu^- \phi^+) - ig\frac{1-2c_w^2}{2c_w}Z_\mu^0(\phi^+ \partial_\mu \phi^- - \phi^- \partial_\mu \phi^+) + \\
 & ig_{sw}A_\mu(\phi^+ \partial_\mu \phi^- - \phi^- \partial_\mu \phi^+) - \frac{1}{4}g^2 W_\mu^+ W_\mu^- [H^2 + (\phi^0)^2 + 2\phi^+ \phi^-] - \\
 & \frac{1}{4}g^2 \frac{1}{c_w^2}Z_\mu^0 Z_\mu^0 [H^2 + (\phi^0)^2 + 2(2s_w^2 - 1)\phi^+ \phi^-] - \frac{1}{2}g^2 \frac{s_w^2}{c_w}Z_\mu^0 \phi^0(W_\mu^+ \phi^- + \\
 & W_\mu^- \phi^+) - \frac{1}{2}ig^2 \frac{s_w^2}{c_w}Z_\mu^0 H(W_\mu^+ \phi^- - W_\mu^- \phi^+) + \frac{1}{2}g^2 s_w A_\mu \phi^0(W_\mu^+ \phi^- + \\
 & W_\mu^- \phi^+) + \frac{1}{2}ig^2 s_w A_\mu H(W_\mu^+ \phi^- - W_\mu^- \phi^+) - g^2 \frac{s_w}{c_w}(2c_w^2 - 1)Z_\mu^0 A_\mu \phi^+ \phi^- - \\
 & g^1 s_w^2 A_\mu A_\mu \phi^+ \phi^- - \bar{e}^\lambda (\gamma \partial + m_e^\lambda) e^\lambda - \bar{\nu}^\lambda \gamma \partial \nu^\lambda - \bar{u}_j^\lambda (\gamma \partial + m_u^\lambda) u_j^\lambda - \\
 & \bar{d}_j^\lambda (\gamma \partial + m_d^\lambda) d_j^\lambda + ig_{sw}A_\mu [-(\bar{e}^\lambda \gamma^\mu e^\lambda) + \frac{2}{3}(\bar{u}_j^\lambda \gamma^\mu u_j^\lambda) - \frac{1}{3}(\bar{d}_j^\lambda \gamma^\mu d_j^\lambda)] + \\
 & \frac{ig}{4c_w}Z_\mu^0 [(\bar{\nu}^\lambda \gamma^\mu (1 + \gamma^5) \nu^\lambda) + (\bar{e}^\lambda \gamma^\mu (4s_w^2 - 1 - \gamma^5) e^\lambda) + (\bar{u}_j^\lambda \gamma^\mu (\frac{4}{3}s_w^2 - \\
 & 1 - \gamma^5) u_j^\lambda) + (\bar{d}_j^\lambda \gamma^\mu (1 - \frac{8}{3}s_w^2 - \gamma^5) d_j^\lambda)] + \frac{ig}{2\sqrt{2}}W_\mu^+ [(\bar{\nu}^\lambda \gamma^\mu (1 + \gamma^5) e^\lambda) + \\
 & (\bar{u}_j^\lambda \gamma^\mu (1 + \gamma^5) C_{\lambda\kappa} d_j^\kappa)] + \frac{ig}{2\sqrt{2}}W_\mu^- [(\bar{e}^\lambda \gamma^\mu (1 + \gamma^5) \nu^\lambda) + (\bar{d}_j^\kappa C_{\lambda\kappa}^\dagger \gamma^\mu (1 + \\
 & \gamma^5) u_j^\lambda)] + \frac{ig}{2\sqrt{2}}\frac{m_e^\lambda}{M}[-\phi^+ (\bar{\nu}^\lambda (1 - \gamma^5) e^\lambda) + \phi^- (\bar{e}^\lambda (1 + \gamma^5) \nu^\lambda)] - \\
 & \frac{g}{2}\frac{m_e^\lambda}{M}[H(\bar{e}^\lambda e^\lambda) + i\phi^0 (\bar{e}^\lambda \gamma^5 e^\lambda)] + \frac{ig}{2M\sqrt{2}}\phi^+ [-m_d^\lambda (\bar{u}_j^\lambda C_{\lambda\kappa} (1 - \gamma^5) d_j^\kappa) + \\
 & m_u^\lambda (\bar{u}_j^\lambda C_{\lambda\kappa} (1 + \gamma^5) d_j^\kappa)] + \frac{ig}{2M\sqrt{2}}\phi^- [m_d^\lambda (\bar{d}_j^\lambda C_{\lambda\kappa}^\dagger (1 + \gamma^5) u_j^\kappa) - m_u^\lambda (\bar{d}_j^\lambda C_{\lambda\kappa}^\dagger (1 - \\
 & \gamma^5) u_j^\kappa) - \frac{g}{2}\frac{m_u^\lambda}{M}H(\bar{u}_j^\lambda u_j^\lambda) - \frac{g}{2}\frac{m_d^\lambda}{M}H(\bar{d}_j^\lambda d_j^\lambda) + \frac{ig}{2}\frac{m_u^\lambda}{M}\phi^0 (\bar{u}_j^\lambda \gamma^5 u_j^\lambda) - \\
 & \frac{ig}{2}\frac{m_d^\lambda}{M}\phi^0 (\bar{d}_j^\lambda \gamma^5 d_j^\lambda)] + \bar{X}^+ (\partial^2 - M^2) X^+ + \bar{X}^- (\partial^2 - M^2) X^- + \bar{X}^0 (\partial^2 - \\
 & \frac{M^2}{c_w^2}) X^0 + \bar{Y} \partial^2 Y + ig_{cw}W_\mu^+ (\partial_\mu \bar{X}^0 X^- - \partial_\mu \bar{X}^+ X^0) + ig_{sw}W_\mu^- (\partial_\mu \bar{Y} X^- - \\
 & \partial_\mu \bar{X}^+ Y) + ig_{cw}W_\mu^- (\partial_\mu \bar{X}^- X^0 - \partial_\mu \bar{X}^0 X^+) + ig_{sw}W_\mu^- (\partial_\mu \bar{X}^- Y - \\
 & \partial_\mu \bar{Y} X^+) + ig_{cw}Z_\mu^0 (\partial_\mu \bar{X}^+ X^+ - \partial_\mu \bar{X}^- X^-) + ig_{sw}A_\mu (\partial_\mu \bar{X}^+ X^+ - \\
 & \partial_\mu \bar{X}^- X^-) - \frac{1}{2}gM[\bar{X}^+ X^+ H + \bar{X}^- X^- H + \frac{1}{c_w^2}\bar{X}^0 X^0 H] + \\
 & \frac{1-2c_w^2}{2c_w}igM[\bar{X}^+ X^0 \phi^+ - \bar{X}^- X^0 \phi^-] + \frac{1}{2c_w}igM[\bar{X}^0 X^- \phi^+ - \bar{X}^0 X^+ \phi^-] + \\
 & igM_{sw}[\bar{X}^0 X^- \phi^+ - \bar{X}^0 X^+ \phi^-] + \frac{1}{2}igM[\bar{X}^+ X^+ \phi^0 - \bar{X}^- X^- \phi^0]
 \end{aligned}$$

$$S = -\frac{1}{16\pi G} \int \sqrt{-g}(R(g)+2\Lambda) d^4x$$



Modified Gravity



Scalar-Tensor Theories

- Field equations:

$$\phi G_{\mu\nu} + \left[\frac{\omega}{2\phi} (\nabla\phi)^2 + V \right] g_{\mu\nu} - \nabla_\mu \nabla_\nu \phi - \frac{\omega}{\phi} \nabla_\mu \phi \nabla_\nu \phi = 8\pi T_{\mu\nu}$$

$$(2\omega + 3) \phi + \omega' (\nabla\phi)^2 + 4V - 2\phi V' = 8\pi T$$

- For Brans-Dicke:

- PPN parameters:

- Newton's constant:

$$\beta_{PPN} = 1, \quad \gamma_{PPN} = \frac{1+\omega}{2+\omega} \Rightarrow \omega \geq 40000$$

$$G = \left(\frac{4+2\omega}{3+2\omega} \right) \frac{1}{\phi} \quad \frac{\dot{G}}{G} \leq 1.7 \cdot 10^{-12} \text{ yr}^{-1}$$

Brans-Dicke Cosmology

- Friedmann-Robertson-Walker metric: $ds^2 = dt^2 - a^2(t)\delta_{ij}dx^i dx^j$

- Friedmann equations:

$$H^2 = \frac{8\pi}{3\phi} \rho_m - H \frac{\dot{\phi}}{\phi} + \frac{\omega}{6} \frac{\dot{\phi}^2}{\phi^2} + \frac{V}{3\phi}$$

$$2\dot{H} + 3H^2 = -\frac{1}{\phi} \left(8\pi p_m + \frac{\omega}{2} \frac{\dot{\phi}^2}{\phi} + 2H\dot{\phi} + \ddot{\phi} \right) + \frac{V}{\phi}$$

- Scalar-field equation:

$$\ddot{\phi} + 3H\dot{\phi} - \frac{8\pi}{2\omega + 3} (\rho_m - 3p_m) = 0 + \frac{2}{2\omega + 3} \left(2V - \phi \frac{dV}{d\phi} \right)$$

- Matter equation: $\dot{\rho}_m + 3H(\rho_m + p_m) = 0$


Dark Energy in Brans-Dicke Cosmology

- Effective Dark Energy sector:


$$\rho_{DE} = \frac{3}{8\pi} \left(-H\dot{\phi} + \frac{\omega}{6} \frac{\dot{\phi}^2}{\phi} \right) + \frac{V}{8\pi}$$

$$\Rightarrow w_{DE} = \frac{p_{DE}}{\rho_{DE}}$$

$$p_{DE} = \frac{1}{8\pi} \left(\frac{\omega}{2} \frac{\dot{\phi}^2}{\phi} + 2H\dot{\phi} + \ddot{\phi} \right) - \frac{V}{8\pi}$$



$$V(\phi) = \frac{V_0}{\phi^2}$$



Scalar-Tensor Theories

- Most general 4D scalar-tensor theories having second-order field equations:

$$L_H = \sum_{i=2}^5 L_i$$

$$L_2[K] = K(\varphi, X)$$

$$L_3[G_3] = -G_3(\varphi, X) \diamond \varphi$$

$$L_4[G_4] = G_4(\varphi, X) R + G_{4,X} \left[(\diamond \varphi)^2 - (\nabla_\mu \nabla_\nu \varphi) (\nabla^\mu \nabla^\nu \varphi) \right]$$

$$L_5[G_5] = G_5(\varphi, X) G_{\mu\nu} (\nabla^\mu \nabla^\nu \varphi) - \frac{1}{6} G_{5,X} \left[(\diamond \varphi)^3 - 3(\diamond \varphi) (\nabla_\mu \nabla_\nu \varphi) (\nabla^\mu \nabla^\nu \varphi) + 2(\nabla^\mu \nabla_\alpha \varphi) (\nabla^\alpha \nabla_\beta \varphi) (\nabla^\beta \nabla_\mu \varphi) \right]$$

$$X = -\partial^\mu \varphi \partial_\mu \varphi / 2$$

[G. Horndeski, Int. J. Theor. Phys. 10
1]

Horndeski Theories

- Most general 4D scalar-tensor theories having second-order field equations:

$$L_H \doteq \sum_{i=2}^5 L_i$$

$$L_2[K] = K(\varphi, X)$$

$$L_3[G_3] = -G_3(\varphi, X) \diamond \varphi$$

$$L_4[G_4] = G_4(\varphi, X) R + G_{4,X} \left[(\diamond \varphi)^2 - (\nabla_\mu \nabla_\nu \varphi) (\nabla^\mu \nabla^\nu \varphi) \right]$$

$$L_5[G_5] = G_5(\varphi, X) G_{\mu\nu} (\nabla^\mu \nabla^\nu \varphi) - \frac{1}{6} G_{5,X} \left[(\diamond \varphi)^3 - 3(\diamond \varphi) (\nabla_\mu \nabla_\nu \varphi) (\nabla^\mu \nabla^\nu \varphi) + 2(\nabla^\mu \nabla_\alpha \varphi) (\nabla^\alpha \nabla_\beta \varphi) (\nabla^\beta \nabla_\mu \varphi) \right]$$

$$X = -\partial^\mu \varphi \partial_\mu \varphi / 2$$

[G. Horndeski, Int. J. Theor. Phys. 10

1



Coincides with Generalized Galileon theories

$$\varphi \rightarrow \varphi + c, \partial_\mu \varphi \rightarrow \partial_\mu \varphi + b_\mu$$

[Nicolis, Rattazzi, Trincherini, PRD 79]

Solving H0 tensions in Horndeski Gravity



$$G_4 = 1/(16\pi G) \text{ and } G_3 = 0, \quad K = -V(\phi) + X$$

$$\rho_{DE} = 2X - K + 2H^3 X \dot{\phi} (5G_{5,X} + 2XG_{5,XX}),$$

$$p_{DE} = K - 2XG_{5,X} (2H^3 \dot{\phi} + 2H\dot{H}\dot{\phi} + 3H^2\ddot{\phi}) - 4H^2X^2\ddot{\phi}G_{5,XX}$$

$$\frac{G_{eff}}{G} = \frac{1}{2} \left(G_4 - 2XG_{4,X} + XG_{5,\phi} - \dot{\phi}HXG_{5,X} \right)^{-1}$$

- Model I: $G_5(X) = \xi X^2$



Solving H0 tensions in Horndeski Gravity

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{\text{Pl}}^2}{2} R + G_2(X) + G_3(X) \square \phi \right]$$

$$G_2(X) = -c_2 M_2^{4(1-p)} (-X/2)^p, \quad G_3(X) = -c_3 M_3^{1-4p_3} (-X/2)^{p_3}$$

$$H_0 = 72_{-5}^{+8} \text{ km s}^{-1} \text{ Mpc}^{-1} \text{ at 95\% CL,}$$

[N. Frusciante, S. Peirone, L. Atayde, A. De Felice, PRD 101]

$$G_2(X) = a_1 X + a_2 X^2, \quad G_3(X) = 3a_3 X$$

$$H_0 = (69.3_{-3.0}^{+3.6}) \text{ km s}^{-1} \text{ Mpc}^{-1} \text{ at 95\%}$$

[S. Peirone, G. Benevento, N. Frusciante, S. Tsujikawa, PRD 100]

Bi-scalar Theories

- Modified gravity propagating 2+2 dof's

$$S = \int d^4x \sqrt{-g} f(R, (\nabla R)^2, \diamond R)$$

- For

$$f(R, (\nabla R)^2, \diamond R) = K(R, (\nabla R)^2) + Q(R, (\nabla R)^2) \diamond R$$

[Naruko, Yoshida, Mukohyama CQG 33]

$$\Rightarrow S = \int d^4x \sqrt{-g} \left[\frac{1}{2} \hat{R} - \frac{1}{2} \hat{g}^{\mu\nu} \partial_\mu \chi \partial_\nu \chi - \frac{1}{\sqrt{6}} e^{-\sqrt{2/3} \chi} \hat{g}^{\mu\nu} Q \partial_\mu \chi \partial_\nu \varphi + \frac{1}{4} e^{-2\sqrt{2/3} \chi} K + \frac{1}{2} e^{-\sqrt{2/3} \chi} Q \diamond \varphi - \frac{1}{4} e^{-\sqrt{2/3} \chi} \varphi \right]$$

$$K = K(\varphi, B), G = G(\varphi, B), B = 2e^{\sqrt{2/3} \chi} g^{\mu\nu} \nabla_\mu \varphi \nabla_\nu \varphi$$

Bi-scalar Theories

- Modified gravity propagating 2+2 dof's

- For

$$f(R, (\nabla R)^2, \diamond R) = K(R, (\nabla R)^2) + Q(R, (\nabla R)^2) \diamond R$$

[Naruko, Yoshida, Mukohyama CQG 33]

$$\Rightarrow S = \int d^4x \sqrt{-g} \left[\frac{1}{2} \hat{R} - \frac{1}{2} \hat{g}^{\mu\nu} \partial_\mu \chi \partial_\nu \chi - \frac{1}{\sqrt{6}} e^{-\sqrt{2/3} \chi} \hat{g}^{\mu\nu} Q \partial_\mu \chi \partial_\nu \varphi + \frac{1}{4} e^{-2\sqrt{2/3} \chi} K + \frac{1}{2} e^{-\sqrt{2/3} \chi} Q \diamond \varphi - \frac{1}{4} e^{-\sqrt{2/3} \chi} \varphi \right]$$



- eg.:

$$K = K(\varphi, B), G = G(\varphi, B), B = 2e^{\sqrt{2/3} \chi} g^{\mu\nu} \nabla_\mu \varphi \nabla_\nu \varphi$$

$$K(\varphi, B) = \frac{\varphi}{2}, G(\varphi, B) = \xi B$$

$$\rho_{DE} = \frac{1}{2} \dot{\chi}^2 - \frac{1}{8} e^{-2\sqrt{2/3} \chi} (1 - 2e^{\sqrt{2/3} \chi}) \varphi - \xi \dot{\varphi}^3 (\sqrt{6} \dot{\chi} - 6H)$$



$$p_{DE} = \frac{1}{2} \dot{\chi}^2 + \frac{1}{8} e^{-2\sqrt{2/3} \chi} (1 - 2e^{\sqrt{2/3} \chi}) \varphi - \frac{1}{3} \xi \dot{\varphi}^2 (\sqrt{6} \dot{\chi} + 6\ddot{\varphi})$$



[Saridakis, Tsoukalas PRD 93]

Solving H0 tensions in Bi-scalar Gravity

- Model I: $\mathcal{K}(\phi, B) = \frac{1}{2}\phi - \frac{\zeta}{2}B$ and $\mathcal{G}(\phi, B) = 0$



[M. Petronikou, E.N.Saridakis, Universe 9]

Running Vacuum

- Upgrade the cosmological constant Λ (vacuum energy) to a **running vacuum**:

$$\begin{aligned} 3H^2 &= 8\pi G(H) (\rho_m + \rho_r + \rho_\Lambda(H)) \\ 3H^2 + 2\dot{H} &= -8\pi G(H) (p_r - \rho_\Lambda(H)) \end{aligned}$$


$$p_{\text{RVM}}^{\text{vac}} = -\rho_{\text{RVM}}^{\text{vac}}$$

$$\rho_\Lambda(H; \nu, \alpha) = \frac{3}{8\pi G} \left(c_0 + \nu H^2 + \frac{2}{3} \alpha \dot{H} \right) + \mathcal{O}(H^4)$$

[Sola, Gomez-Valent, Perez *Astrophys. J.*
836]

[Basilakos, Polarski, Sola *PRD*86]

Solving the tensions in Running Vacuum



[J. Sola, A. Gomez-Valent, J. de Cruz Perez, C. MorenoPulido
CQG 37]

“Those that do not know geometry are not allowed to enter”.

Front Door of Plato's Academy



Descriptions of Gravity

- Einstein 1916: **General Relativity:**
energy-momentum source of spacetime
Curvature
Levi-Civita connection: Zero Torsion
- Einstein 1928: **Teleparallel Equivalent of GR:**
Weitzenböck connection: Zero Curvature

[Cai, Capozziello, De Laurentis, Saridakis,
Rept.Prog.Phys. 79]


$$\left\{ \begin{smallmatrix} \alpha \\ \mu\nu \end{smallmatrix} \right\} = \frac{1}{2} g^{\alpha\lambda} (g_{\lambda\nu,\mu} + g_{\mu\lambda,\nu} - g_{\mu\nu,\lambda}). \quad (1.3)$$

The corresponding covariant derivative will be denoted by $\mathcal{D}$ so that we will have $\mathcal{D}_\alpha g_{\mu\nu} = 0$. A general connection $\Gamma^\alpha_{\mu\nu}$ then admits the following convenient decomposition:


$$\Gamma^\alpha_{\mu\nu} = \left\{ \begin{smallmatrix} \alpha \\ \mu\nu \end{smallmatrix} \right\} + K^\alpha_{\mu\nu} + L^\alpha_{\mu\nu} \quad (1.4)$$

with


$$K^\alpha_{\mu\nu} = \frac{1}{2} T^\alpha_{\mu\nu} + T_{(\mu}{}^\alpha{}_{\nu)}, \quad L^\alpha_{\mu\nu} = \frac{1}{2} Q^\alpha_{\mu\nu} - Q_{(\mu}{}^\alpha{}_{\nu)} \quad (1.5)$$



The rotation of a vector transported along a closed curve is given by the curvature: General Relativity.



The non-closure of parallelograms formed when two vectors are transported along each other is given by the torsion: Teleparallel Equivalent of General Relativity.



The variation of the length of a vector as it is transported is given by the non-metricity: Symmetric Teleparallel Equivalent of General Relativity.

Metric-Affine Modified Gravity




FIG. 1. Subclasses of metric-affine geometry, depending on the properties of connection.

$$S_{\text{GR}} = \frac{1}{2\kappa^2} \int \left\{ g^{\mu\nu} \hat{R}_{\mu\nu} + \lambda_{(1)}^{\mu\nu\lambda} T_{\mu\nu\lambda} + \lambda_{(2)}^{\mu\nu\lambda} Q_{\mu\nu\lambda} \right\} \sqrt{-g} d^4x,$$

$$S_{\text{total}} = S_{\text{GR}} + S_{\text{matter}},$$

Teleparallel Equivalent of General Relativity (TEGR)

- Let's start from the **simplest torsion-based** gravity formulation, namely **TEGR**:
- Vierbeins** e_A^μ : four linearly independent fields in the **tangent space**

$$g_{\mu\nu}(x) = \eta_{AB} e_\mu^A(x) e_\nu^B(x)$$
- Use **curvature-less Weitzenböck connection** instead of **torsion-less Levi-Civita** one: $\Gamma_{\nu\mu}^\lambda = e_A^\lambda \partial_\mu e_\nu^A$
- Torsion tensor**:

$$T_{\mu\nu}^\lambda = \Gamma_{\nu\mu}^\lambda - \Gamma_{\mu\nu}^\lambda = e_A^\lambda (\partial_\mu e_\nu^A - \partial_\nu e_\mu^A)$$
- Lagrangian** (imposing coordinate, Lorentz, parity invariance, and up to 2nd order in torsion tensor)

$$L \equiv T = \frac{1}{4} T^{\rho\mu\nu} T_{\rho\mu\nu} + \frac{1}{2} T^{\rho\mu\nu} T_{\nu\mu\rho} - T_{\rho\mu}^\rho T_\nu^{\nu\mu}$$

- Completely equivalent**
with
GR at the level of
equations

f(T) Gravity and f(T) Cosmology

- **f(T) Gravity**: Simplest torsion-based modified gravity
- Generalize T to **f(T)** (inspired by **f(R)**)

$$S = \frac{1}{16\pi G} \int d^4x e [T + f(T)] + S_m$$

- Equations of motion:

$$e^{-1} \partial_\mu (e e_A^\rho S_\rho^{\mu\nu}) (1 + f_T) - e_A^\lambda T_{\mu\lambda}^\rho S_\rho^{\nu\mu} + e_A^\rho S_\rho^{\mu\nu} \partial_\mu (T) f_{TT} - \frac{1}{4} e_A^\nu [T + f(T)] = 4\pi G e_A^\rho T_\rho^\nu \{EM\}$$

f(T) Gravity and f(T) Cosmology

- **f(T) Gravity**: Simplest torsion-based modified gravity
- Generalize T to **f(T)** (inspired by **f(R)**)

$$S = \frac{1}{16\pi G} \int d^4x e [T + f(T)] + S_m$$

- Equations of motion:

$$e^{-1} \partial_\mu (e e_A^\rho S_\rho^{\mu\nu}) (1 + f_T) - e_A^\lambda T_{\mu\lambda}^\rho S_\rho^{\nu\mu} + e_A^\rho S_\rho^{\mu\nu} \partial_\mu (T) f_{TT} - \frac{1}{4} e_A^\nu [T + f(T)] = 4\pi G e_A^\rho T_\rho^\nu \{EM\}$$

- **f(T) Cosmology**: Apply in FRW geometry:

$$e_\mu^A = \text{diag}(1, a, a, a) \Rightarrow ds^2 = dt^2 - a^2(t) \delta_{ij} dx^i dx^j \text{ (not unique choice)}$$

- Friedmann equations:

$$H^2 = \frac{8\pi G}{3} \rho_m - \frac{f(T)}{6} - 2f_T H^2$$

$$\dot{H} = - \frac{4\pi G(\rho_m + p_m)}{1 + f_T - 12H^2 f_{TT}}$$

- Find easily

$$T = -6H^2$$

f(T) Cosmology: Background

- Effective **Dark Energy** sector:

$$\rho_{DE} \equiv \frac{3}{8\pi G_N} \left[-\frac{f}{6} + \frac{T f_T}{3} \right],$$

$$P_{DE} \equiv \frac{1}{16\pi G_N} \left[\frac{f - f_T T + 2T^2 f_{TT}}{1 + f_T + 2T f_{TT}} \right]$$

$$w_{DE} = -\frac{f - T f_T + 2T^2 f_{TT}}{[1 + f_T + 2T f_{TT}][f - 2T f_T]}$$

[Linder PRD 82]

- Interesting cosmological behavior: **Late-time acceleration**, **Inflation** etc

[Cai, Capozziello, De Laurentis, Saridakis, Rept.Prog.Phys. 79]

f(T) Cosmology: Perturbations

- For **scalar perturbations**:

$$e_{\mu}^0 = \delta_{\mu}^0 (1 + \psi), e_{\mu}^{\alpha} = \delta_{\mu}^{\alpha} \alpha (1 - \varphi)$$

$$\Rightarrow ds^2 = (1 + 2\psi) dt^2 - a^2 (1 - 2\varphi) \delta_{ij} dx^i dx^j$$

- Obtain **Perturbation Equations**.

$$\ddot{\delta} + 2H\dot{\delta} - 4\pi G_{\text{eff}} \rho \delta \approx 0$$

$$Q(a) = \frac{G_{\text{eff}}(a)}{G_N} = \frac{1}{1 + f_T}$$

[Chen, Dent, Dutta, Saridakis PRD 83],

[Dent, Dutta, Saridakis JCAP 1101]

$$\begin{aligned} \delta T_0^0 &= -\delta\rho_m \\ \delta T_0^i &= a^{-2}(\rho_m + p_m)(-\partial_i \delta u) \\ \delta T_i^0 &= (\rho_m + p_m)(\partial_i \delta u) \\ \delta T_i^j &= \delta_{ij}\delta p_m + \partial_i \partial_j \pi^S. \end{aligned}$$

$$\begin{aligned} E_0 &\equiv (1 + f'_0)(\nabla^2 \phi) + 6(1 + f'_0)H\dot{\phi} \\ &\quad + 6(1 + f'_0)H^2\psi - 3f'_1 H^2 \\ &\quad - \frac{T_1 + f_1}{4} = -4\pi G\delta\rho_m, \end{aligned}$$

$$\begin{aligned} E_0^i &\equiv (1 + f'_0)\partial_i \dot{\phi} + (1 + f'_0)H\partial_i \psi \\ &\quad - 12H\dot{H}f_0''\partial_i \phi = -4\pi G(\rho_m + p_m)\partial_i \delta u, \\ E_a^0 &\equiv 12H^2\partial_i \delta_a^i (\dot{\phi} + H\psi)f_0'' - (1 + f'_0)\partial_i \delta_a^i (\dot{\phi} + H\psi) \\ &\quad = 4\pi G(\rho_m + p_m)\partial_i \delta_a^i \delta u, \end{aligned}$$

$$\begin{aligned} E_a^i \delta_i^a &\equiv \frac{f_1}{a} (-3H^2 - \dot{H}) + \frac{f_1''}{a} (12H^2 \dot{H}) \\ &\quad - \frac{(1 + f'_0)}{2a} \sum_{b \neq a} \partial^j \delta_j^b \partial_i \delta_b^i (\psi - \phi) \\ &\quad - \frac{\phi(T_0 + f_0)}{4a} - \frac{T_1 + f_1}{4a} \\ &\quad + \frac{(1 + f'_0)}{a} [6H\dot{\phi} + 6H^2\psi - 3H^2\dot{\phi} \\ &\quad \quad + \ddot{\phi} + \dot{H}(2\psi - \phi) + H\dot{\psi}] \\ &\quad + \frac{f_0''}{a} (-24H\dot{H}\dot{\phi} - 48\psi H^2\dot{H} - 12H^2\ddot{\phi} \\ &\quad \quad - 12H^3\dot{\psi} + 12H^2\dot{H}\phi) \\ &\quad = \frac{4\pi G}{a} (p_m \phi + \delta p_m), \end{aligned}$$

$$\begin{aligned} E_{b; b \neq a}^i \delta_i^a &\equiv \frac{(1 + f'_0)}{2} \partial_j \delta_b^j \partial_i^a \delta_i^a (\phi - \psi) \\ &\quad = 4\pi G a^2 \partial_j \delta_b^j \partial_i^a \delta_i^a \pi^S \end{aligned}$$

Efficient model independent requirements to solve the tensions

We consider a correction in the first Friedmann equation of the form

$$H(z) = -\frac{d(z)}{4} + \sqrt{\frac{d^2(z)}{16} + H_{\Lambda\text{CDM}}^2(z)}, \quad (2)$$

where $H_{\Lambda\text{CDM}}(z) \equiv H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$ is the Hubble rate in ΛCDM , with $\Omega_m = \rho_m/(3M_p^2 H^2)$ the matter density parameter and primes denote derivatives with respect to z .

- If $d < 0$ and is suitably chosen, one can have $H(z \rightarrow z_{\text{CMB}}) \approx H_{\Lambda\text{CDM}}(z \rightarrow z_{\text{CMB}})$ but $H(z \rightarrow 0) > H_{\Lambda\text{CDM}}(z \rightarrow 0)$; i.e., the H_0 tension is solved [one should choose $|d(z)| < H(z)$, and thus, since $H(z)$ decreases for smaller z , the deviation from ΛCDM will be significant only at low redshift].
- Since the friction term in (1) increases, the growth of structure gets damped, and therefore, the σ_8 tension is also solved.

Solving H_0 and S8 tensions in $f(T)$ Gravity

- We consider the following ansatz:

$$f(T) = -[T + 6H_0^2(1 - \Omega_{m0}) + F(T)], \quad (9)$$

where $F(T)$ describes the deviation from GR

The first Friedmann equation becomes

$$T(z) + 2\frac{F'(z)}{T'(z)}T(z) - F(z) = 6H_{\Lambda\text{CDM}}^2(z). \quad (10)$$

- In order to solve the H_0 tension, we need $T(0) = 6H_0^2 \simeq 6(H_0^{\text{CC}})^2$, with $H_0^{\text{CC}} = 74.03 \text{ km s}^{-1} \text{ Mpc}^{-1}$, while in the early era of $z \gtrsim 1100$ we require the Universe expansion to evolve as in ΛCDM , namely $H(z \gtrsim 1100) \simeq H_{\Lambda\text{CDM}}(z \gtrsim 1100)$. This implies $F(z)|_{z \gtrsim 1100} \simeq cT^{1/2}(z)$ (the value $c = 0$ corresponds to standard GR, while for $c \neq 0$ we obtain ΛCDM too).

Solving H0 and S8 tensions in f(T) Gravity

The effective gravitational coupling is given by

$$G_{\text{eff}} = \frac{G_N}{1 + F_T} . \quad (11)$$

Therefore, the perturbation equation becomes



$$\delta'' + \left[\frac{T'(z)}{2T(z)} - \frac{1}{1+z} \right] \delta' = \frac{9H_0^2 \Omega_{m0}(1+z)}{[1 + F'(z)/T'(z)]T(z)} \delta . \quad (12)$$

Since around the last scattering moment $z \gtrsim 1100$ the Universe should be matter-dominated, we impose

$\delta'(z)|_{z \gtrsim 1100} \simeq -\frac{1}{1+z} \delta(z)$, while at late times we look for $\delta(z)$ that leads to an $f\sigma_8$ in agreement with redshift survey observations.

Solving H0 and S8 tensions in f(T) Gravity



By solving (10) and (12) with initial and boundary conditions at $z \sim 0$ and $z \sim 1100$, we can find the functional forms for the free functions of the $f(T)$ gravity that we consider, namely, $T(z)$ and $F(z)$, that can alleviate both H_0 and σ_8 tensions.





$$\text{Model-1: } F(T) \approx 375.47 \left(\frac{T}{6H_0^2} \right)^{-1.65}$$

$$\text{Model-2: } F(T) \approx 375.47 \left(\frac{T}{6H_0^2} \right)^{-1.65} + 25 T^{1/2}.$$

Solving H0 and S8 tensions in f(T) Gravity



Parameter	CMB + BAO	CMB + BAO + H_0
$10^2 \omega_b$	$2.235^{+0.013}_{-0.013}$	$2.235^{+0.013}_{-0.013}$
ω_{cdm}	$0.1181^{+0.001}_{-0.001}$	$0.118^{+0.001}_{-0.001}$
$100\theta_s$	$1.041^{+0.00027}_{-0.00027}$	$1.041^{+0.00030}_{-0.00027}$
$\ln 10^{10} A_s$	$3.078^{+0.023}_{-0.023}$	$3.08^{+0.022}_{-0.022}$
n_s	$0.9678^{+0.0039}_{-0.0039}$	$0.9684^{+0.0039}_{-0.0044}$
τ_{reio}	$0.073^{+0.012}_{-0.013}$	$0.075^{+0.012}_{-0.012}$
n	$0.0043^{+0.0033}_{-0.0039}$	$0.0054^{+0.0020}_{-0.0020}$
$\log \alpha$	$10.00^{+0.081}_{-0.12}$	$10.03^{+0.06}_{-0.06}$
Ω_{F0}	$0.73^{+0.021}_{-0.028}$	$0.738^{+0.015}_{-0.015}$
H_0	$72.4^{+3.3}_{-4.1}$	$73.5^{+2.1}_{-2.1}$
σ_8	$0.855^{+0.023}_{-0.033}$	$0.866^{+0.02}_{-0.02}$
$\chi^2_{min}/2$	6480.48	6482.27




[S-F Yan, P. Zhang, J-W Chen, X_Z Zhang, Y-F Cai, E.N. Saridakis, PRD 101]

[J-W Chen, W. Luo, Y-F Cai, E.N. Saridakis, PRD 102]

[S. Basilakos, S. Nesseris, F. Anagnostopoulos, E.N. Saridakis, JCAP 2019]

Viabie f(T) models



[Nesseris, Basilakos, Saridakis, Perivolaropoulos, PRD 88]

In other modified gravities: Not possible

- This behavior **is not possible** in other **modified gravities**. e.g.:


$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2} f(R, \phi, X) + \mathcal{L}_m \right) \quad X = -g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi$$

$$G_{\text{eff}}(a, k)/G_{\text{N}} = \frac{1}{F} \frac{f_{,X} + 4 \left(f_{,X} \frac{k^2}{a^2} \frac{F_{,R}}{F} + \frac{F_{,\phi}^2}{F} \right)}{f_{,X} + 3 \left(f_{,X} \frac{k^2}{a^2} \frac{F_{,R}}{F} + \frac{F_{,\phi}^2}{F} \right)} \quad F = F(R, \phi, X) = \partial_R f(R, \phi, X)$$

- $G_{\text{eff}}/G_{\text{N}} > 1$ for all models that **do not have ghosts** (i.e. with $f_R, f_{RR} > 0$).

- On the contrary, **f(T) gravity** has **second-order field equations** and moreover **perturbations are stable** in a large part of the parameter phase.

f(Q) gravity



affine connection $\Gamma^{\alpha}_{\mu\nu}$ can be decomposed as

$$\Gamma^{\alpha}_{\mu\nu} = \hat{\Gamma}^{\alpha}_{\mu\nu} + K^{\alpha}_{\mu\nu} + L^{\alpha}_{\mu\nu}, \quad (1)$$

where $\hat{\Gamma}^{\alpha}_{\mu\nu}$ is the Levi-Civita connection,

$$K^{\alpha}_{\mu\nu} = \frac{1}{2}T^{\alpha}_{\mu\nu} + T^{\alpha}_{(\mu \nu)} \quad (2)$$

is the contortion tensor with $T^{\alpha}_{\mu\nu}$ the torsion tensor, and

$$L^{\alpha}_{\mu\nu} = \frac{1}{2}Q^{\alpha}_{\mu\nu} - Q^{\alpha}_{(\mu \nu)} \quad (3)$$

is the disformation tensor arising from the non-metricity

$$Q_{\alpha\mu\nu} \equiv \nabla_{\alpha}g_{\mu\nu}, \quad (4)$$

FIG. 1. Subclasses of metric-affine geometry, depending on the properties of connection.

f(Q) gravity

$$T^\lambda_{\mu\nu} \equiv \Gamma^\lambda_{\mu\nu} - \Gamma^\lambda_{\nu\mu}$$

$$R^\sigma_{\rho\mu\nu} \equiv \partial_\mu \Gamma^\sigma_{\nu\rho} - \partial_\nu \Gamma^\sigma_{\mu\rho} + \Gamma^\alpha_{\nu\rho} \Gamma^\sigma_{\mu\alpha} - \Gamma^\alpha_{\mu\rho} \Gamma^\sigma_{\nu\alpha} \quad (5)$$

while the nonmetricity can be expressed as

$$Q_{\rho\mu\nu} \equiv \nabla_\rho g_{\mu\nu} = \partial_\rho g_{\mu\nu} - \Gamma^\beta_{\rho\mu} g_{\beta\nu} - \Gamma^\beta_{\rho\nu} g_{\mu\beta} . \quad (6)$$

$$Q = -\frac{1}{4}Q_{\alpha\beta\gamma}Q^{\alpha\beta\gamma} + \frac{1}{2}Q_{\alpha\beta\gamma}Q^{\gamma\beta\alpha} + \frac{1}{4}Q_\alpha Q^\alpha - \frac{1}{2}Q_\alpha \tilde{Q}^\alpha , \quad (7)$$

where $Q_\alpha \equiv Q_\alpha^\mu{}_\mu$, and $\tilde{Q}^\alpha \equiv Q_\mu^{\mu\alpha}$.

f(Q) gravity

$$S = -\frac{1}{2} \int d^4x \sqrt{-g} f(Q). \quad (8)$$

$$\begin{aligned} & \frac{2}{\sqrt{-g}} \nabla_\alpha \left\{ \sqrt{-g} g_{\beta\nu} f_Q \left[-\frac{1}{2} L^{\alpha\mu\beta} + \frac{1}{4} g^{\mu\beta} (Q^\alpha - \tilde{Q}^\alpha) \right. \right. \\ & \quad \left. \left. - \frac{1}{8} (g^{\alpha\mu} Q^\beta + g^{\alpha\beta} Q^\mu) \right] \right\} \\ & + f_Q \left[-\frac{1}{2} L^{\mu\alpha\beta} - \frac{1}{8} (g^{\mu\alpha} Q^\beta + g^{\mu\beta} Q^\alpha) \right. \\ & \quad \left. + \frac{1}{4} g^{\alpha\beta} (Q^\mu - \tilde{Q}^\mu) \right] Q_{\nu\alpha\beta} + \frac{1}{2} \delta_\nu^\mu f = T^\mu_\nu, \quad (9) \end{aligned}$$

with $f_Q = \partial f / \partial Q$.

f(Q) cosmology

- Background:

$$6f_Q H^2 - \frac{1}{2}f = \rho_m,$$

$$(12H^2 f_{QQ} + f_Q)\dot{H} = -\frac{1}{2}(\rho_m + p_m). \quad (11)$$

$$Q = 6H^2, \quad (12)$$

f(Q) cosmology

■ Perturbations:

$$-a^2 \delta \rho = 6 (f_Q + 12a^{-2} \mathcal{H}^2 f_{QQ}) \mathcal{H} (\mathcal{H} \phi + \varphi') + 2f_Q k^2 \psi - 2 [f_Q + 3a^{-2} f_{QQ} (\mathcal{H}' + \mathcal{H}^2)] \mathcal{H} k^2 B. \quad (19)$$

$$\begin{aligned} \frac{1}{2} a^2 (\rho + p) v = & [f_Q + 3a^{-2} f_{QQ} (\mathcal{H}' + \mathcal{H}^2)] \mathcal{H} \phi \\ & + 6a^{-2} f_{QQ} \mathcal{H}^2 \varphi' - 9a^{-2} f_{QQ} (\mathcal{H}' - \mathcal{H}^2) \mathcal{H} \varphi \\ & + f_Q \psi' - a^{-2} f_{QQ} \mathcal{H}^2 k^2 B, \end{aligned} \quad (20)$$

$$\begin{aligned} \frac{1}{2} a^2 \delta p = & (f_Q + 12a^{-2} f_{QQ} \mathcal{H}^2) (\mathcal{H} \phi' + \varphi'') + \left[f_Q \left(\mathcal{H}' + 2\mathcal{H}^2 - \frac{1}{3} k^2 \right) + 12a^{-2} f_{QQ} \mathcal{H}^2 (4\mathcal{H}' - \mathcal{H}^2) + 12a^{-2} \frac{df_{QQ}}{d\tau} \mathcal{H}^3 \right] \phi \\ & + 2 \left[f_Q + 6a^{-2} f_{QQ} (3\mathcal{H}' - \mathcal{H}^2) + 6a^{-2} \frac{df_{QQ}}{d\tau} \mathcal{H} \right] \mathcal{H} \varphi' + \frac{1}{3} f_Q k^2 \psi \\ & - \frac{1}{3} (f_Q + 6a^{-2} f_{QQ} \mathcal{H}^2) k^2 B' - \frac{1}{3} \left[2f_Q + 3a^{-2} f_{QQ} (5\mathcal{H} - \mathcal{H}^2) + 6a^{-2} \frac{df_{QQ}}{d\tau} \mathcal{H} \right] \mathcal{H} k^2 B, \end{aligned} \quad (21)$$

f(Q) cosmology

■ Perturbations:

$$\delta' = (1 + w) (-k^2 v - k^2 B + 3\varphi') + 3\mathcal{H} \left(w\rho - \frac{\delta p}{\rho} \right), \quad (22)$$


$$v' = -\mathcal{H} (1 - c_s^2) v + \frac{\delta p}{\rho + p} + \phi. \quad (23)$$

$$\begin{aligned} & - f_{QQ} \mathcal{H} [2\mathcal{H}\varphi' + (\mathcal{H}' + \mathcal{H}^2) \phi + (\mathcal{H}' - \mathcal{H}^2) (\psi - B')] \\ & - \left[f_{QQ} \left(\mathcal{H}'^2 + \mathcal{H}\mathcal{H}'' - 3\mathcal{H}^2\mathcal{H}' - \frac{1}{3}\mathcal{H}^2 k^2 \right) \right. \\ & \left. + \frac{df_{QQ}}{d\tau} (\mathcal{H}' - \mathcal{H}^2) \mathcal{H} \right] B = 0, \end{aligned} \quad (24)$$


$$\delta'' + \mathcal{H}\delta' = \frac{4\pi G\rho}{f_Q} \delta, \quad (30)$$

$$G_{eff} \equiv \frac{G}{f_Q}, \quad (31)$$

Solving the tensions in $f(Q)$ gravity



[F. Anagnostopoulos, S. Basilakos, E.N.Saridakis, JCAP 2019]



[S. Peirone, G. Benevento, N. Frusciante, S. Tsujikawa, PRD 100]

Conclusions

- i) **Astrophysics** and **Cosmology** have become **precision** sciences.
- ii) A **huge amount of accumulating data** suggest possible **tensions** with **theoretical predictions** of Λ CDM paradigm.
- iii) **New Physics** or **paradigm shift** may be the **way out**
- iv) We can **modify** the **Universe content**, the **interactions**, or/and the **gravitational theory**. Historically, modified gravity has been proven to be the solution quite often.



Conclusions

- i) **Astrophysics** and **Cosmology** have become **precision** sciences.
- ii) A **huge amount of accumulating data** suggest possible **tensions** with **theoretical predictions** of Λ CDM paradigm.
- iii) **New Physics** or **paradigm shift** may be the **way out**
- iv) We can **modify** the **Universe content**, the **interactions**, or/and the **gravitational theory**. Historically, modified gravity has been proven to be the solution quite often.



THANK YOU!